

Choline

2565

Please Copy for Your Patients

Choline Helps Your Body Metabolize Fats and Support Nervous System Function

Choline is an important 'vitamin-like' substance that makes up a large portion of all cell membranes and is necessary for the body to transport different fats in the bloodstream. It is especially known for its ability to help keep the liver free from fatty buildup. Choline is also a key component of two important elements of the nervous system, acetylcholine and sphingomyelin, both necessary for the normal functioning of the nervous system.†

How Choline Keeps You Healthy

Supports healthy nervous system function

Choline and calcium are needed to transmit nerve impulses from the brain throughout the central nervous system. Choline enhances brain function and memory.†

Keeps your heart and liver healthy

Fat is automatically transported out of the liver during the same process in which choline becomes lecithin. The primary function of choline is to metabolize fats throughout the body.†

Maintains cellular health

Choline furnishes an essential structural component of many biological membranes. Choline donates methyl groups that are necessary in order to synthesize other important compounds needed for various physiological processes.†



Introduced in:

1947

Content:

90 Tablets

Supplement Facts:

Serving Size: 1 tablet
Servings per Container: 90

		%DV
Calories	2	
Choline	180 mg	

Choline 2565



800-558-8740 • www.standardprocess.com

† These statements have not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Choline

What Makes Choline Unique

Unique Product Attributes

This is a vegetarian product

A supplement form of choline provides consistent bioavailability

- Choline levels vary considerably in food, making it difficult to determine the amount of choline actually available for utilization by the body†

Unique Processing

Degreed microbiologists and chemists in our on-site laboratories constantly conduct bacterial and analytical tests on raw materials, product batches, and finished products

- Ensures consistent quality and safety

Vitamin and mineral analyses validate product content and specifications

- Assures high-quality essential nutrients are delivered

Whole Food Philosophy

Dr. Lee challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists—known and unknown—bioactivity is markedly enhanced over synthetic nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to a synthetic or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

Excellent source of choline.

Ingredients: Choline bitartrate, honey, and calcium stearate.

Suggested Use: One tablet per meal, or as directed.

Sold to health care professionals.

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food ingredients. See the supplement facts for Choline.

Anderson L. E. 1998. *Mosby's Medical, Nursing, & Allied Health Dictionary*. 5th ed. Mosby; St. Louis: 246, 326.
Balch J.F., Balch P.A. 1997. *Prescription for Nutritional Healing*. 2nd ed. Avery Publishing Group: Garden City Park, 17, 23, 55.
Berdanier C.D. 1995. *Advanced Nutrition Micronutrients*. CRC Press: Boca Raton. 130-133, 158-172.
Blusztajn J.K., et al. Imprinting of hippocampal metabolism of choline by its availability during gestation: implications for cholinergic neurotransmission. *Paris Journal of Physiology*. Jun-Aug 1998; 92(3-4): 199-203.
Caamano J., et al. Effects of CDP choline on cognition and cerebral hemodynamics in patients with Alzheimer's disease. *Methods Find Exp Clin Pharmacol*. Apr 1994; 16(3): 211-218.
Corona G.L., et al. Clinical and biochemical responses to therapy in Alzheimer's disease and multi-infarct dementia. *Eur Arch Psychiatry Neurol Sci*. 1989; 239(2): 79-86.
Darvesh S., et al. Cholinesterases in cardiac ganglia and modulation of canine intrinsic cardiac neuronal activity. *Journal of the Autonomic Nervous System*. Jul 15 1998; 71(2-3): 75-84.
Denda A., et al. Inhibition by piroxicam of oxidative DNA damage, liver cirrhosis and development of enzyme-altered nodules caused by a choline-deficient, L-amino acid-defined diet in rats. *Carcinogenesis*. Oct 1997; 18(10): 1921-1930.
Denda A., et al. Prevention by inhibitors of arachidonic acid cascade of liver carcinogenesis, cirrhosis and oxidative DNA damage caused by a choline-deficient, L-amino acid-defined diet in rats. *Mutation Research*. Jun 18 1998; 402(1-2): 279-288.
Emmert J.J., et al. Hepatic betaine-homocysteine methyltransferase activity in the chicken is influenced by dietary intake of sulfur amino acids, choline and betaine. *Journal of Nutrition*. Aug 1996; 126(8): 2050-2058.
Geiger J., et al. The effect increased choline levels on the synthesis and release of acetylcholine in heart atria in white rats. *Bratisl Lek Listy*. Dec 1992; 93(12): 610-614.
Ghoshal A.K. New insight into the biochemical pathology of liver in choline deficiency. *Critical Review of Biochemical Molecular Biology*. 1995; 30(4): 263-273.
Giovannini M.G., et al. Effect of subchronic treatment with metrifonate and tacrine on brain cholinergic function in aged F344 rats. *European Journal of Pharmacology*. Jul 31 1998; 354(1): 17-24.
Hacker H.J., et al. Pyruvate kinase isoenzyme shift from L1-type to M2-type is a late event in hepatocarcinogenesis induced in rats by a choline-deficient/DL-ethionine-supplemented diet. *Carcinogenesis*. Jan 1998; 19(1): 99-107.
Holmes McNary M.Q., et al. Choline and choline esters in human and rat milk and in infant formulas. *American Journal of Clinical Nutrition*. Oct 1996; 64(4): 572-56.
Kenney J.L., Carlberg K.A. The effect of choline and myo-inositol on liver and carcass fat levels in aerobically trained rats. *International Journal of Sports Medicine*. Feb 1995; 16(2): 114-116.
Kobayashi E., et al. Inhibitory effects of N,N'-diphenyl-p-phenylenediamine on the early stage of the enhanced hepatocarcinogenesis caused by coadministration of ethionine and a choline deficient L-amino acid defined diet in rats. *Exp Toxicol Pathol*. Jun 1996; 48(4): 275-282.
Lindmar R., et al. Characterization of choline efflux from the perfused heart at rest and after muscarinic receptor activation. *Naunyn-Schmiedberg Arch Pharmacol*. Mar 1986; 332(3): 224-229.
Machlin L.J. 1984. *Handbook of Vitamins*. Marcel Dekker, Inc.: New York. 550-561.

Mock T., et al. Effects of fatty acids on phosphatidylcholine biosynthesis in isolated hamster heart. *Biochemical Cellular Biology*. May 1986; 64(5): 413-417.
Nakae D., et al. Production of both 8-hydroxydeoxyguanosine in liver DNA and gamma-glutamyltransferase-positive hepatocellular lesions in rats given a choline-deficient, L-amino acid-defined diet. *Japanese Journal of Cancer Research*. Nov 1990; 81(11): 1081-1084.
Pieri C. Chronic choline treatment improves the "in vivo" membrane permeability of old hepatocytes to Rb+. *Boll Soc Ital Biol Sper*. Oct 1989; 65(10): 909-915.
Pitchford P. 1993. *Healing with Whole Foods*. North Atlantic Books: Berkeley. 122, 177-185.
Rabkin S.W. Effects of chlorpromazine and trifluoperazine on choline metabolism and phosphatidylcholine biosynthesis in cultured chick heart cells under normoxic and anoxic conditions. *Biochemical Pharmacology*. Jul 15 1989; 38(14): 2349.
Rabkin S.W. Effect of exogenous CDP-choline on choline metabolism in isolated adult rat ventricular myocytes under normoxic and hypoxic conditions. *Cellular Biochemical Function*. Jun 1993; 11(2): 137-143.
Sandmann J., et al. The effects of phorbol esters on choline phospholipid hydrolysis in heart and brain. *European Journal of Pharmacology*. Mar 13 1990; 188(2-3): 89-95.
Sardesai V.M. 1998. *Introduction to Clinical Nutrition*. Marcel Dekker, Inc.: New York. 79-84, 233-235.
Scheider W.L. 1983. *Nutrition, Basic Concepts and Applications*. McGraw Hill Book Company: New York. 202-203, 222-224.
Seca E., et al. Choline feeding depresses the phospholipase C activity in the regenerating liver of female rats. *Boll Soc Ital Biol Sper*. Nov-Dec 1996; 72(11-12): 325-329.
Seca E., et al. The delay in rat liver regeneration by choline is associated to alteration in C-myc expression. *Boll Soc Ital Biol Sper*. Jul-Aug 1996; 72(7-8): 217-222.
Sheard N.F., et al. Plasma choline concentration in humans fed parenterally. *American Journal of Clinical Nutrition*. Feb 1986; 43(2): 219-224.
Sheard N.F., Zeisel S.H. Choline: an essential dietary nutrient? *Nutrition*. Jan-Feb 1989; 5(1): 1-5.
Shils M.E., Young V.R. 1988. *Modern Nutrition in Health and Disease*. 7th ed. Lea & Febiger: Philadelphia. 142-154, 440-449.
Shinozuka H., et al. Choline deficiency and chemical carcinogenesis. *Adv Exp Med Biol*. 1986; 206, 253-267.
Sugiyama T., et al. Enhancing effect of a choline-deficient diet on alterations of hepatic drug-metabolizing enzymes in hepatitis- and hepatoma-predisposed rats (LEC rats). *Japanese Journal of Cancer Research*. Apr 1991; 82(4): 390-396.
Tamura K., et al. Inhibition by N-(4-hydroxyphenyl) retinamide and all-trans-retinoic acid of exogenous and endogenous development of putative preneoplastic, glutathione S-transferase placental form-positive lesions in the livers of rats. *Carcinogenesis*. Nov 1997; 18(11): 2133-2141.
Tsujiuchi T. Increased telomerase activity in hyperplastic nodules and hepatocellular carcinomas induced by a choline-deficient L-amino acid-defined diet in rats. *Japanese Journal of Cancer Research*. Nov 1996; 87(11): 1111-1115.
Tver D.F., Percy R. 1989. *The Nutrition and Health Encyclopedia*. 2nd ed. Van Nostrand Reinhold. New York. 117-118, 86-88, 266.
Unelius L., et al. Choline+ is a low-affinity ligand for alpha-1-adrenoceptors. *Journal of Biochemical Pharmacology*. Oct 1994; 48(7): 1519-1523.
West Sutor C.J., Crowley F., Merrily R.N. 1984. *Nutrition, Principles and Application in Health Promotion*. 2nd ed. J.B. Lippincott Company: Philadelphia. 49, 65-66.
Wilson E.D., Fisher K.H., Fuqua M.E. 1965. *Principles of Nutrition*. 2nd ed. John Wiley & Sons: New York. 136-150, 308-309.
Yokota K., et al. Effects of a choline-deficient diet and a hypolipidemic agent on single glutathione S-transferase placental form-positive hepatocytes in rat liver. *Japanese Journal of Cancer Research*. Feb 1990; 81(2): 129-134.